

TO OPEN...
CUT HERE!



32070

Hi-Yield®



IMPROVED SLUG & SNAIL BAIT

Controlled Release Pellet to Control Slugs and Snails Around Listed Vegetables, Citrus, and Ornamentals

- Attractants Draw In All Species Of Slugs and Snails
- Long Lasting. The All Weather Formula is effective up to 3 to 4 weeks
- Use Around Listed Fruits, Vegetables and Ornamentals
- Controls Giant African Land Snail
- Easy Spread Mini Pellets
- 2.5 Lbs covers 11,000 sq. ft.



ACTIVE INGREDIENT:
Metaldehyde 3.25%

OTHER INGREDIENTS:..... 96.75%

TOTAL:..... 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

This pesticide may be harmful to children and fatal to domestic animals if ingested.

Exclude children and pets from treated areas until the applied product is no longer visible.

For additional precautionary measures, see "Directions for Use" and "Storage and Disposal".



32070

NET WEIGHT 2.5 LBS. (1.13 KG)

Hi-Yield®

IMPROVED SLUG & SNAIL BAIT

32070

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful if inhaled. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

IMPORTANT: This product may be harmful to children and fatal to domestic animals if ingested. Children and dogs may be attracted to Metaldehyde products and their packaging. Keep children and pets out of treated areas from the start of application until the applied product is no longer visible.

ENVIRONMENTAL HAZARDS

This product is toxic to birds, mammals, and mollusks (e.g. Clams, Oysters, Scallops, Mussels). Do not apply product near water (including garden ponds) or storm drains. Do not apply if heavy rain is expected. Apply this product only around gardens and ornamental plants. Do not contaminate water when disposing of equipment or container wash water. Collect and remove any product that is spilled.

PRODUCT INFORMATION

HI-YIELD® IMPROVED SLUG & SNAIL BAIT is a soil surface treatment for control of Snails and Slugs around listed vegetable crops and ornamental gardens. When applied as directed, **HI-YIELD® IMPROVED SLUG & SNAIL BAIT** will withstand rain and sprinkling and will offer protection to gardens for several weeks.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label. Use in accordance with Precautionary Statements and Directions For Use. Not for use on plants being grown for sales or other commercial use, or for commercial seed production, or research purposes.

KEEP OUT OF REACH OF CHILDREN

Hi-Yield®

IMPROVED

SLUG & SNAIL BAIT

FIRST AID

| | |
|-------------------------------|--|
| If Swallowed | <ul style="list-style-type: none"> • Call Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by the Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person. |
| If Inhaled | <ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a Poison Control Center or doctor for further treatment advice. |
| If In Eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice. |
| If On Skin Or Clothing | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a Poison Control Center or doctor for treatment advice. |

POISONING HOTLINE

Seek medical care as soon as possible after exposure. For incidents involving animals, seek veterinary care as soon as possible after exposure. Have the product container or label with you when you call a Poison Control Center or doctor or veterinarian, or when going for treatment. For information on this pesticide product (including health concerns and pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

METALDEHYDE IS A SYSTEMIC TOXIN. THERE IS NO ANTIDOTE. THE FOLLOWING INSTRUCTIONS ARE NOT INTENDED FOR LAYPEOPLE. THERE IS NO HOME TREATMENT. EXPOSED INDIVIDUALS AND ANIMALS SHOULD RECEIVE EMERGENCY CARE AS QUICKLY AS POSSIBLE.

Note to Physician: Dust inhalation is usual route of accidental exposure; the product is rarely ingested. Treatment is symptomatic and supportive. Monitor severe exposures closely: seizures, electrolyte imbalances, acidosis, dehydration, and increased body temperature have been reported.

Note to Veterinarian: Animals typically are exposed via ingestion. Treatment is symptomatic and supportive. If significant ingestion is suspected, and the animal is asymptomatic, consider induction of emesis or gastric lavage and administration of activated charcoal, cathartic and enema to limit absorption. Monitor asymptomatic and symptomatic animals closely. Treat seizures, electrolyte imbalances, acidosis, dehydration, and increased body temperature. Monitor liver function post recovery.

**In the event of a medical or chemical emergency, contact
North America 1-800-255-3924 or Worldwide Intl. + 01- 813-248-0585**

PRECAUTIONS AND RESTRICTIONS

-This pesticide may be harmful to children and fatal to dogs and other domestic animals if ingested. Children and dogs may be attracted to Metaldehyde products, both in the package and when applied. Children and domestic animals must be kept out of treated areas from the start of application until the applied product is no longer visible.

-Application of this Metaldehyde product is prohibited unless children and domestic animals can be excluded from the treated areas from the start of application until the applied product is no longer visible.

-Applications to turf and Dichondra are prohibited. Application is permitted only to non-turf areas (i.e., on soil or mulch) directly surrounding plants listed for use on label.

-Broadcast and foliar applications of this product are prohibited. **Application is permitted only as a barrier around gardens or individual plants. Application is prohibited within one foot of vegetable plants or fruit-bearing plants not listed on this label.**

- Avoid applying pellets in piles that may attract pets.
- Certain cultural practices may reduce damage from Snails and Slugs. These practices include: eliminating hiding places by removing leaf litter and debris in the garden; removing Slugs and Snails during daylight hours from nearby leaf litter, mulch, and other protected areas; watering in the morning to reduce humidity during the night when Snails and Slugs are active.

- Do not apply to dry soil. Make applications following irrigation.

- Do not re-water for 48 hours after application.

- Apply any time of day, but evening applications are preferred.

- Do not reuse measuring utensils for food use.

- This product may be harmful to Daylilies and Clematis. Do not use on these plants.

APPLICATION INFORMATION

Apply pellets as a barrier around plants by distributing pellets evenly in a line at the following rates, pellets per linear foot.

| Crop Use Site | Maximum Pellets per Linear Foot | Minimum Days Between Retreatment | Maximum Number of Applications Per Growing Season | Preharvest Interval | Maximum number of applications per year |
|--|---------------------------------|----------------------------------|---|---------------------|---|
| Artichoke -perennial -annual | 20 20 | 18 18 | 6 2 | 0 0 | 12 4 |
| Berries: Caneberries (bingleberry, black raspberry, blackberry, boysenberry, dewberry, loganberry, lowberry, marionberry, olallieberry, red raspberry, wild raspberry, youngberry), Bushberries (aronia berry, blueberry, buffalo currant, Chilean guava, black currant, red currant, elderberry, European barberry, gooseberry, cranberry, honeysuckle, huckleberry, jostaberry, juneberry, lingonberry, salal, sea buckthorn), Berry-low growing (bearberry, bilberry, blueberry, cloudberry, muntries, partridgeberry) | 18 | 14 | 3 | 0 | 3 |
| Strawberries | 20 | 14 | 3 | 0 | 6 |
| Celery, Rhubarb, Swiss Chard, Chinese Celery, Celtuce, Cardoon, and Florence Fennel | 20 | 21 | 4 | 1 | 8 |
| Peas and Bean | 20 | 7 | 3 | Pre-bloom | 6 |
| Prickly Pear Cactus | 18 | 30 | 3 | 0 | 3 |
| Tomatoes | 20 | 14 | 3 | 0 | 6 |
| Lettuce, Cole Crops and the following Leafy Greens: broccoli, Brussels Sprouts, cabbage, cauliflower, cavalo, broccolo, collards, kale, kohlrabi, mizuna, mustard greens, spinach, rape greens | 20 | 14 | 3 | 0 | 6 |
| Ornamental Plants | 20 | 21 | 6 | 0 | 6 |

632070

Manufactured For:




230 FM 87 • BONHAM, TEXAS 75418

EPA Reg. No. 71096-7-7401

EPA Est. No. 7401-TX-1

Product Questions? 855-270-4776



Hi-Yield[®]

IMPROVED
**SLUG & SNAIL
BAIT**
32070

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.
PESTICIDE STORAGE: This product can be harmful to children and fatal to dogs and other domestic animals if ingested. Dogs have been known to ingest Metaldehyde after opening or tearing packaging. Store this product in its original packaging, in a cool, secure location, and out of the reach of children and pets.

CONTAINER DISPOSAL: If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If Partly Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.